

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

EA

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>A61K 9/16, 9/24, 9/26, 9/30, 9/48, 9/52</b>		A1	(11) International Publication Number: <b>WO 99/47123</b>
			(43) International Publication Date: 23 September 1999 (23.09.99)

(21) International Application Number: <b>PCT/US99/05111</b>	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: <b>9 March 1999 (09.03.99)</b>	Published <i>With international search report.</i>
(30) Priority Data: <b>09/040,794 18 March 1998 (18.03.98) US</b>	
(71) Applicant: <b>BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).</b>	
(72) Inventors: <b>ULLAH, Ismat; 2 Mockingbird Court, Cranbury, NJ 08512 (US). JAIN, Nemichand, B.; 19 Abbington Lane, Cranbury, NJ 08512 (US).</b>	
(74) Agents: <b>RODNEY, Burton et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).</b>	

(54) Title: **PHARMACEUTICAL COMPOSITION CONTAINING A STATIN AND ASPIRIN**

## (57) Abstract

A pharmaceutical composition is provided which is useful for cholesterol lowering and reducing the risk of a myocardial infarction, which includes a statin, such as pravastatin, lovastatin, simvastatin, atorvastatin, cerivastatin or fluvastatin, in combination with aspirin, in a manner to minimize interaction of aspirin with the statin and minimize side effects of aspirin. A method for lowering cholesterol and reducing risk of a myocardial infarction employing such composition is also provided.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PHARMACEUTICAL COMPOSITION CONTAINING A STATIN AND ASPIRIN

Field of the Invention

5        The present invention relates to a pharmaceutical composition which includes a statin cholesterol lowering agent and aspirin in a manner to minimize interaction of aspirin with the statin, for use in lowering cholesterol and reducing risk of a myocardial infarction, and to a  
10      method for lowering cholesterol and reducing risk of a myocardial infarction employing such composition.

Background of the Invention

15      The use of aspirin for reducing the risk of a myocardial infarction and the use of statins for lowering cholesterol and preventing or treating atherosclerosis and cardiovascular disease and cerebrovascular disease are well documented. In fact, it is not uncommon that patients having elevated cholesterol levels who are at high risk for  
20      a myocardial infarction take both a statin and aspirin. However, use of both a statin and aspirin may require special care to insure that drug interaction, including physical and chemical incompatibility, and side effects, are kept to a minimum while achieving maximum benefit from  
25      these drugs.

With regard to possible drug interaction, aspirin is an acid, while some of the statins, such as pravastatin, atorvastatin and cerivastatin, are alkali salts. Thus, mixing of such statins (alkali salts) with aspirin could  
30      result in aspirin hydrolysis as well as statin degradation. Pravastatin, on the other hand, is also a very acid labile compound. When pravastatin and aspirin are combined, the aspirin could cause pravastatin degradation which could result in lower bioavailability of pravastatin.

35      Aspirin is known for causing gastrointestinal bleeding when used for long-term therapy. It is therefore

desirable in long-term aspirin therapy that the aspirin be provided in a form which minimizes side effects.

In view of the above, it is seen that there is a long-felt want in patients required to take both a statin and aspirin for a statin-aspirin formulation which provides for maximum cholesterol lowering and reduction of risk of a myocardial infarction without the undesirable side effects and drug interaction normally associated with use of such combination.

10

#### Description of the Invention

In accordance with the present invention, a pharmaceutical composition is provided which includes a statin cholesterol lowering agent and aspirin, which provides for maximum patient benefits including maximum cholesterol lowering and reduced risk of a myocardial infarction with minimal physical and chemical incompatibility (including minimal statin:aspirin interaction), and reduced side effects normally associated with use of such drugs.

In addition, in accordance with the present invention, a method is provided for lowering serum cholesterol, preventing or inhibiting or treating atherosclerosis, and/or reducing risk of or treating a cardiovascular event or disease including coronary artery disease and cerebrovascular disease, wherein a pharmaceutical composition containing a combination of a statin cholesterol lowering agent and aspirin in a single dosage form, in a manner so as to minimize interaction of the statin and aspirin, is administered to a patient in need of treatment.

Preferred pharmaceutical compositions of the present invention may take the form of several different embodiments. Thus, in one embodiment of the present invention, a pharmaceutical composition is provided wherein the statin (including any statin cholesterol lowering agent) and aspirin are formulated together in a single

tablet. The tablet of the invention is preferably in the form of a bilayered tablet which includes a first layer and a second layer. Aspirin, in the form of granules of preselected size will be present in the first layer 5 together with optional excipients as described hereinafter, while the statin will be present in the second layer which optionally may include one or more buffering agents (as necessary to prevent undesirable statin/aspirin interaction) and optionally one or more excipients as 10 described hereinafter.

In addition, the bilayered tablet of the invention may include an outer protective coating or finishing layer as described hereinafter.

Another embodiment of the present invention 15 comprises a cored tablet which includes a core and a buffering layer or outer coat which can be compressed onto the core as a dry coat. The core will preferably include compressed aspirin granules while the buffering layer or outer coat will include a statin (including any statin 20 cholesterol lowering agent) together with one or more buffering agents and optional excipients.

Provision of aspirin in the core and statin in the buffering layer will effectively reduce the aspirin side effects and also minimize drug incompatibilities while 25 providing maximum efficacy.

The so-described cored tablet may also optionally include an outer protective coating or finishing layer as described hereinafter.

In addition, in accordance with the present 30 invention, a pharmaceutical composition is provided which is in the form of a tablet or capsule which includes a mixture of aspirin granules having an enteric coating and particles or granules of a statin. Such a combination will provide maximum efficacy while minimizing side effects 35 resulting from prolonged aspirin therapy.

In the above embodiment containing enteric coated aspirin, the statin may include any statin cholesterol

lowering agent, but preferably is simvastatin, lovastatin or cerivastatin.

In yet another embodiment of the pharmaceutical composition of the present invention, enteric coated aspirin granules as described above may be further coated with a protective coating or finishing layer. The double coated particles of aspirin can be mixed with any statins such as pravastatin, atorvastatin, simvastatin, lovastatin, and cerivastatin powders or granules, and the mixture can be encapsulated or tableted as described herein. In such case, further coating of the enteric coated aspirin particles is desired to minimize interaction of alkaline pravastatin, atorvastatin or cerivastatin with enteric coated aspirin. This combination will protect the integrity of the enteric coat and minimize the side effects normally resulting from prolonged aspirin therapy. The aspirin and the statin granules do not need to be mixed together; these can even be encapsulated separately into the same capsule shells in two shots.

Another embodiment of the pharmaceutical composition of the invention includes granules of enteric coated aspirin and enteric coated statin (including any statin cholesterol lowering agent), in the same dosage form such as compressed tablets or capsules.

The tablets containing the enteric coated granules of aspirin and statin may also include an outer protective coating or finishing layer.

In a further embodiment of the pharmaceutical composition of the invention, where aspirin side effects are not an issue, for example, where low dose aspirin is present (81 mg or less), the composition of the invention may comprise a mixture of aspirin granules and statin (including any statin cholesterol lowering agent, preferably, simvastatin, lovastatin or enteric coated particles of pravastatin or particles of pravastatin, atorvastatin and cerivastatin containing an outer protective coating or finishing layer); the above mixture

may take the form of compressed tablets or capsules (where the mixture can be encapsulated separately in two shots in the same capsule shells).

5

Detailed Description of the Invention

The pharmaceutical composition of the invention which includes a combination of a statin and aspirin is effective in preventing, reducing and/or treating elevated cholesterol levels (such as in hypercholesterolemia), 10 atherosclerosis, cardiovascular events and disease including coronary events and cerebrovascular events, and coronary artery disease and/or cerebrovascular disease.

The terms "cardiovascular event(s)" and "cardiovascular disease" as employed herein refer to 15 coronary and/or cerebrovascular event(s) and disease including primary myocardial infarction, secondary myocardial infarction, myocardial ischemia, angina pectoris (including unstable angina), congestive heart failure, sudden cardiac death, cerebral infarction, cerebral 20 thrombosis, cerebral ischemia, transient ischemic attack and the like.

The term "coronary artery disease" (CAD) as employed herein refers to diseases including atherosclerosis of the coronary arteries, previous myocardial infarction, 25 ischemia, angina pectoris and/or heart failure.

The term "cerebrovascular disease" as employed herein refers to diseases including atherosclerosis of the intracranial and/or extracranial arteries, cerebral infarction, cerebral thrombosis, cerebral ischemia, stroke, 30 and/or transient ischemic attacks.

Aspirin will preferably be employed in the form of salicylic acid acetate also referred to as acetylsalicylic acid.

The pharmaceutical composition of the invention in 35 the form of a tablet or capsule will include aspirin in amounts from about 10 to about 800 mg, preferably 50 to about 650 mg.

The aspirin for use in forming the pharmaceutical composition of the invention will preferably be in the form of granules having an average particle size within the range from about 10  $\mu\text{m}$  to about 2 mm, more preferably from 5 about 0.25 mm to about 1.0 mm.

Statin cholesterol lowering agents suitable for use herein will include HMG CoA reductase inhibitors such as pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin and other statins such as 10 fluindostatin and preferably pravastatin, simvastatin, atorvastatin or cerivastatin.

The pharmaceutical composition of the invention will contain a statin such as pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin in 15 an amount as normally employed for such statin as exemplified in the 52nd edition of the Physician's Desk Reference (PDR) (1998). Thus, depending upon the particular statin, it may be employed in amounts within the range from about 0.1 mg to 2000 mg per day in single or 20 divided doses, and preferably from about 0.2 to about 200 mg per day. Most preferably for pravastatin, a daily dosage of 10 to 40 mg may be employed; for lovastatin, a daily dosage of 10 to 80 mg may be employed, for simvastatin a daily dosage of 5 to 40 mg may be employed; 25 for atorvastatin, a daily dosage of 10 to 80 mg may be employed, for fluvastatin, a daily dosage of 20 to 80 mg may be employed; and for cerivastatin, a daily dosage of 0.2-0.3 mg may be employed.

In forming the pharmaceutical composition of the 30 invention in the form of a bilayered tablet, the first layer containing aspirin will also preferably include bulking agents such as lactose, microcrystalline cellulose, wood cellulose, corn starch, modified corn starch, calcium phosphate, sugar, dextrose, mannitol or sorbitol. The 35 bulking agent will be present in an amount from about 1 to about 90%, preferably from about 5 to about 85% by weight of the first layer containing aspirin.

The first layer may also include a tabletting lubricant, such as zinc stearate, magnesium stearate, calcium stearate, talc, carnauba wax, stearic acid, palmitic acid or hydrogenated vegetable oils and fats, in 5 an amount within the range from about 0.01 to about 4%, and preferably 0.02 to about 2% by weight of the first layer.

The second layer of the bilayered tablet containing statin cholesterol lowering agent will usually include a bulking agent such as lactose, microcrystalline cellulose, 10 modified corn starch, calcium phosphate or other bulking agent as set out above for the first layer, in an amount within the range from about 1 to about 90%, preferably from about 5 to about 85% by weight of the second layer. In addition, the second layer may include a binder such as 15 corn starch, pregelatinized starch, polyvinyl pyrrolidone (PVP), hydroxypropylmethyl cellulose (HPMC), ethyl cellulose, cellulose acetate and the like, in an amount within the range from about 0.5 to about 20%, preferably from about 1 to about 10% by weight of the second layer, 20 and a tabletting lubricant such as magnesium stearate, zinc stearate, or other lubricant as set out above with respect to the first layer in an amount from about 0.01 to about 4%, preferably from about 0.02 to about 2% by weight of the second layer.

25 The buffering agents present in the second layer may include conventional acid buffers such as calcium carbonate, magnesium oxide, magnesium carbonate, magnesium hydroxide, aluminum hydroxide, dihydroxyaluminum sodium carbonate, aluminum magnesium hydroxide sulfate or aluminum 30 hydroxide magnesium carbonate co-dried gel, or mixtures of one or more thereof, in amounts as needed to insure that the aspirin will be sufficiently buffered to inhibit GI side effects. Thus, amounts of buffering agent within the range from about 10 to about 1000 mg, preferably from about 35 50 to about 500 mg will be employed depending upon the amount of aspirin present in the first layer.

In forming the bilayered tablet of the invention, the first layer containing aspirin may be prepared by conventional wet granulation or dry granulation (compaction) techniques.

5 The second layer containing statin and buffers may be prepared by conventional wet granulation or dry granulation (compaction) techniques.

10 The first and second layers may then be compressed and combined to form a bilayered tablet employing conventional bilayer tabletting equipment.

15 Other conventional ingredients which may optionally be present in either of the two layers include preservatives, stabilizers, anti-adherents or silica flow conditioners or glidants, such as Syloid brand silicon dioxide as well as antioxidants such as Vitamin E, Vitamin C, and folic acid, Vitamin B<sub>6</sub> and Vitamin B<sub>12</sub>.

20 The bilayer tablet of the invention may also include an outer protective coating layer which may comprise from 0 to about 15% by weight of the bilayer tablet. The outer protective coating layer which is applied over the bilayered tablet may comprise any conventional coating formulations and will include one or more film-formers or binders, such as a hydrophilic polymer like hydroxypropylmethyl cellulose (HPMC) and a hydrophobic polymer like ethyl cellulose, cellulose acetate, polyvinyl alcohol-maleic anhydride copolymers, acrylic copolymers,  $\beta$ -pinene polymers, glyceryl esters of wood resins and the like, and one or more plasticizers, such as polyethylene glycol, triethyl citrate, diethyl phthalate, propylene glycol, 30 glycerin, butyl phthalate, castor oil and the like.

35 The film formers are applied from a solvent system containing one or more solvents including water, alcohols like methyl alcohol, ethyl alcohol or isopropyl alcohol, ketones like acetone, or ethylmethyl ketone, chlorinated hydrocarbons like methylene chloride, dichloroethane, and 1,1,1-trichloroethane.

The pharmaceutical composition of the invention in the form of a cored tablet wherein the aspirin forms the core, and statin plus buffering agent are present in a surrounding coat layer, may be prepared employing 5 conventional cored tablet technology. Thus, the aspirin containing core (including excipients and other ingredients as described for the first layer in the bilayered tablet of the invention) may be formed in a manner similar to the first layer of the bilayered tablet as described 10 hereinbefore. The buffering layer containing statin as well as excipients and other ingredients (as described hereinbefore for the second layer of the bilayered tablet of the invention) may be compressed onto the core as a dry coat.

15 The so-formed cored tablet may be coated with an outer protective coating layer as described above for the bilayered tablet.

Another embodiment of the pharmaceutical composition of the invention is formed of tablets or capsules 20 containing a mixture of enteric coated aspirin granules, and a statin such as lovastatin, simvastatin or cerivastatin, which may be in the form of a tablet or capsule.

The aspirin granules can be coated with conventional 25 enteric polymers coatings in aqueous or non-aqueous systems. For example, Eudragit L-30D-55 (acrylic acid copolymers-Rohm Pharma) (5 to 25% solids) containing 10 to 15% of diethylphthalate (w/w) as plasticizer can be used in an aqueous system.

30 Other conventional enteric polymer coating systems may be employed such as Eudragit R and S series resins, (acrylic acid copolymers-Rohm Pharma), cellulose acetate phthalate, cellulose acetate maleate, cellulose acetate succinate, hydroxypropylmethyl cellulose phthalate, 35 hydroxypropylmethylcellulose acetate succinate, and the like, and a suitable plasticizer such as triethyl citrate, diethyl phthalate, tributyl citrate, triacetin, dibutyl

phthalate dibutyl sebicate, Myvacet 940, and other commonly used plasticizers as may be suitable for particular enteric polymers can be used. It will be appreciated that any polymer with suitable plasticizer can be used in aqueous or 5 non-aqueous system to form an enteric coating on the aspirin granule or particle.

In another embodiment of the pharmaceutical composition of the invention, the enteric coated aspirin granules described above may be further coated with an 10 outer protective finishing coat or layer as described hereinbefore.

The double coated aspirin granules can be mixed with a statin such as pravastatin, atorvastatin, simvastatin, lovastatin, fluvastatin or cerivastatin powders or granules 15 and the mixture can be encapsulated or tableted as described above.

In yet another embodiment of the pharmaceutical composition of the invention, aspirin is enteric coated as described above and the statin (pravastatin, simvastatin, 20 lovastatin, atorvastatin, fluvastatin and cerivastatin) can optionally be enteric coated. The statins can be coated in the form of pure drugs or after spheronization or agglomeration. The particles for coating do not need to be perfectly spherical. These could be rods or irregular 25 particles. The enteric coated particles of the two drugs (aspirin and statin) can be tableted or encapsulated together. As described above, appropriate excipients (fillers, binders, disintegrants, and lubricant, etc.) can be used to facilitate tabletting. This statin:aspirin 30 combination will minimize side effects of aspirin, and eliminate chemical incompatibility.

If, aspirin side effects are not an issue, especially at lower (e.g., 80 mg) aspirin dosages, then aspirin granules (including uncoated aspirin) can be mixed 35 with simvastatin, lovastatin and fluvastatin powder or granules for tabletting or for encapsulating.

In yet another embodiment, aspirin granules can be mixed with enteric coated particles of pravastatin, cerivastatin and atorvastatin and the mixture can be tableted or encapsulated or the two granules can be 5 encapsulated in two shots in the same capsule shells.

In carrying out the method of the present invention, the pharmaceutical composition of the invention containing the combination of the statin cholesterol lowering drug and aspirin may be administered to mammalian species, such as 10 monkeys, dogs, cats, rats, humans, etc., and, as described hereinbefore, may be incorporated in a tablet or capsule. The above dosage forms will also include the necessary carrier material, excipient, lubricant, buffer, 15 antibacterial, bulking agent (such as mannitol), anti-oxidants such as Vitamin C and Vitamin E, as well as Vitamin B<sub>6</sub>, Vitamin B<sub>12</sub>, folic acid, sodium bisulfite, and the like.

The dose administered must be adjusted according to age, weight and condition of the patient, as well as the 20 route of administration, dosage form and regimen and the desired result.

The compositions described above may be administered in the dosage forms as described above in single or divided doses of one to four times daily. It may be advisable to 25 start a patient on a low dose combination and work up gradually to a high dose combination.

Tablets of various sizes can be prepared, e.g., of about 2 to 2000 mg in total weight, containing the active substances in the ranges described above, with the 30 remainder being a physiologically acceptable carrier of other materials according to accepted pharmaceutical practice. These tablets can, of course, be scored to provide for fractional doses in some cases. Gelatin capsules can be similarly formulated.

Liquid formulations can also be prepared by dissolving or suspending one or the combination of active substances in a conventional liquid vehicle acceptable for pharmaceutical administration so as to provide the desired 5 dosage in one to four teaspoonsful.

Such dosage forms can be administered to the patient on a regimen of one to four doses per day.

In general, formulating the compositions, as described herein, the active substances, in the amounts 10 described above, are compounded as described herein (according to accepted pharmaceutical practice) with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in the particular type of unit dosage form.

15 Illustrative of the excipients which may be incorporated in tablets are the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; an excipient such as dicalcium phosphate or cellulose; a disintegrating agent such as corn starch, potato starch, 20 alginic acid, sodium starch glycolate or the like; a lubricant such as stearic acid, zinc stearate or magnesium stearate; a sweetening agent such as sucrose, aspartame, lactose or saccharin; a flavoring agent such as orange, peppermint, oil of wintergreen or cherry. When the dosage 25 unit form is a capsule, it may contain in addition to materials of the above type a liquid carrier such as a fatty oil. As indicated, various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets or capsules 30 may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds, water, alcohol or the like as the carrier, glycerol as solubilizer, sucrose as sweetening agent, methyl and propyl parabens as preservatives, a dye and a flavoring such as cherry or 35 orange.

Some of the active substances described above form commonly known, pharmaceutically acceptable salts such as alkali metal and other common basic salts or acid addition salts, etc. References to the base substances are 5 therefore intended to include those common salts known to be substantially equivalent to the parent compound.

The formulations as described above will be administered for a prolonged period, that is, for as long as the potential for cardiovascular events and disease 10 including coronary artery disease and/or cerebrovascular disease remains or the symptoms continue. Sustained release forms of such formulations which may provide such amounts daily, biweekly, weekly, monthly and the like may also be employed. A dosing period of at least 10 days are 15 required to achieve minimal benefit.

The following Examples represent preferred embodiments of the present invention.

Formulations suitable for oral administration are prepared as described below.

20

Example 1

A bilayered tablet containing aspirin in a first layer and pravastatin in a second layer as described below may be prepared as follows.

25

**General Formula:**

		Amount or %
		<u>in First Layer</u>
<u>First Layer:</u>		
Aspirin granulation		80 mg - 325 mg
30 Lactose/microcrystalline		qs
Cellulose granulation*		
Zinc Stearate		0.1% - 0.5

	Amount in Second Layer
<u>Second Layer:</u>	
Calcium Carbonate	50 mg - 250 mg
Magnesium Oxide	50 mg - 100 mg
5 Magnesium Carbonate	25 mg - 50 mg
Corn Starch	25 mg - 50 mg
Pravastatin	20 mg - 40 mg
Magnesium stearate	0.2% - 0.5%
10 *This is an inert granulation just for the purpose of bulking, if necessary. This will contain 50%-90% lactose anhydrous, 10%-50% microcrystalline cellulose, and 0.1%-0.5% zinc stearate. These ingredients are blended, and appropriate size granules are prepared by conventional dry granulation process. (This being just an inert granulation, any other excipient can be used to prepare granules for bulking by dry or wet granulation processes, so that the granules do not have alkalizing agent and also do not contain excessive moisture and are compatible with 20 aspirin granules. These bulking granules must have enough compatibility to allow compression of two layer tablets).	
15	
20	

**Procedure:**

The aspirin granulation in the first layer is blended with sufficient quantity of the lactose/microcrystalline cellulose granulation as necessary to bulk up in order to have sufficient granulation to compress a satisfactory layer. The aspirin granules along with the bulking granules are blended with zinc stearate as a lubricant. Zinc stearate can be replaced with other non-alkaline lubricants, i.e., Lubritab® or other high melting point hydrogenated powdered waxes.

5      Ingredients in the second layer are wet granulated using starch paste or other wet granulating materials, for example, PVP or HPMC, or can be dry granulated by compaction. The granules can be sized and lubricated. The two tablet layers are compressed using appropriate conventional tools and a suitable bilayer tabletting press, to form the bilayered tablet of the invention.

10     The quantity of the buffering agents used in the second layer can be adjusted as necessary to minimize gastrointestinal side effects. It should be understood that these buffering agents can be replaced with other suitable buffering agents, if desired.

15     The so-formed bilayered tablets may be coated with HPMC (hydroxypropylmethylcellulose) or commercially available Opadry® clear or Dri Klear® (HPMC) or any of these with any desired color. This coat is not limited to HPMC based coats only. Polymers, i.e., Eudragit E30D (acrylic acid copolymer) and others can also be used to give the tablets a finishing coat.

20

**Coating Formula (example):**

Opadry® clear	10% - 30%
Purified water	qs

25     **Procedure:**

30     Opadry® is dispersed in water to prepare a dispersion of 10%-30% solids\*. This dispersion is used for coating the above tablets using conventional coating equipment. The coating of 0.2%-2% or any desired level (based on the weight of the finished coated bilayered tablet) can be applied to the bilayered tablet employing conventional techniques.

35     The so-formed tablets provide maximum benefits while minimizing drug interaction and other undesirable side effects.

It will be understood that pravastatin contained in the buffered layer of the bilayered tablet of the invention may be replaced with equivalent cholesterol lowering amounts of simvastatin, lovastatin, atorvastatin, 5 cerivastatin or fluvastatin.

\*Antifoam emulsion at a level of 0.1 to 2% of solids, can also be included in the formulation.

10 Example 2

Tablets or capsules containing enteric coated aspirin and a statin, which preferably is simvastatin, lovastatin or cerivastatin, having the following composition are prepared as described below.

15

**General Formula:**

Aspirin particles	80 mg - 325 mg
Eudragit L-30D-55	qs
Diethyl Phthalate	qs
20 Statins (simvastatin, lovastatin, or cerivastatin)	Desired Dose (as per PDR)

**Procedure:**

Aspirin particles are coated with enteric polymers 25 in aqueous or non-aqueous systems. Eudragit L-30D-55 containing 10%-15% of diethyl phthalate (w/w) is used in an aqueous system. The coating suspension is prepared having solid contents of 10%-30%.

To prepare the coating suspension, diethyl phthalate 30 is added to the Eudragit L-30D-55 and the contents stirred till diethyl phthalate is completely dissolved. This is diluted with water to obtain the suspension with desired solid contents. Using this enteric coating suspension, the aspirin particles are coated in a fluid bed coating system 35 using a Wurster insert or with top spray coating, so that aspirin particles of enteric quality can be produced. The enteric coated particles are mixed with statin powders or

granules and the mixtures are encapsulated or tableted using appropriate excipients (fillers, binder, disintegrants, and lubricants). Any of the listed statin can be selected at its desired dose level along with the 5 desired aspirin dose.

The statins can also be granulated, and the statin granules and the enteric coated aspirin granules can be filled separately into the same capsule shell. Statin granules can be prepared by dry or wet granulation 10 processes, using suitable conventional excipients as is well known in the pharmaceutical field.

The above formulations provide maximum benefit while minimizing undesirable side effects and incompatibilities.

15 Example 3

A cored tablet containing an aspirin core and a buffered coating thereon containing a statin having the following composition is prepared as described below.

20 **General Formula:**

	Amount or %
	<u>in Core Layer</u>
<u>Core Layer:</u>	
Aspirin granulation	80 mg - 325 mg
Lactose/microcrystalline	qs
25 Cellulose granulation*	
Zinc Stearate	0.1% - 0.5
	<u>Amount in Second</u>
	<u>Layer</u>
<u>Outer Layer:</u>	
30 Calcium Carbonate	50 mg - 250 mg
Magnesium Oxide	50 mg - 100 mg
Magnesium Carbonate	25 mg - 50 mg
Corn Starch	25 mg - 50 mg
Pravastatin	20 mg - 30 mg
35 Magnesium stearate	0.2% - 0.5%
Filler/Binder**	qs

\*This is an inert granulation just for the purpose of bulking, if necessary. This will contain 50%-90% lactose anhydrous, 10%-50% microcrystalline cellulose, and 0.1%-0.5% zinc stearate. These ingredients are blended, and 5 appropriate size granules are prepared by conventional dry granulation process. (This being just an inert granulation, any other excipient can be used to prepare granules for bulking by dry or wet granulation processes, so that the granules do not have alkalizing agent and also 10 do not contain excessive moisture and are compatible with aspirin granules. These bulking granules must have enough compatibility to allow compression of two layer tablets).

\*\*The Filler/Binder may be any known fillers or tablet 15 binders, such as lactose, microcrystalline cellulose, modified starch, calcium phosphate and the like.

**Procedure:**

The aspirin granulation for the core is blended with 20 sufficient quantity of the lactose/microcrystalline cellulose granulation as necessary to bulk up in order to have sufficient granulation to compress a satisfactory core. The aspirin granules along with the bulking granules are blended with zinc stearate as a lubricant. Zinc 25 stearate can be replaced with other non-alkaline lubricants, i.e., Lubritab® or other high melting point hydrogenated powdered waxes.

Ingredients for the outer layer are wet granulated using starch paste or other wet granulating materials, for 30 example, PVP or HPMC, or can be dry granulated by compaction. The granules can be sized and lubricated. The dry coated tablets can be compressed using appropriate tools and a suitable dry coating tabletting press.

The quantity of the buffering agents used in the 35 outer layer can be adjusted as in Example 1. Other known buffering agents may be used as well.

What is Claimed is:

1. A pharmaceutical composition comprising a statin cholesterol lowering agent and aspirin in a formulation designed to minimize statin:aspirin interaction.
- 5 2. The pharmaceutical composition as defined in Claim 1 wherein the statin and aspirin are formulated together in the same dosage formulation.
- 10 3. The pharmaceutical composition as defined in Claim 2 wherein the statin and aspirin are formulated together in a single tablet.
- 15 4. The pharmaceutical composition as defined in Claim 3 wherein the tablet is a bilayered tablet wherein the aspirin is present in a first layer, and the statin is present in a second layer.
- 20 5. The pharmaceutical composition as defined in Claim 4 wherein the layer containing the statin also includes one or more buffering agents.
- 25 6. The pharmaceutical composition as defined in Claim 3 wherein the tablet includes a core and a coating layer surrounding said core and wherein one of the statin and aspirin is present in the core and the other is present in the coating layer surrounding the core.
- 30 7. The pharmaceutical composition as defined in Claim 6 wherein the aspirin is present in the core and the statin is present in the coating layer.
- 35 8. The pharmaceutical composition as defined in Claim 7 wherein the coating layer also includes one or more buffering agents.
9. The pharmaceutical composition as defined in Claim 1 wherein the statin is pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin or cerivastatin.
10. The pharmaceutical composition as defined in Claim 1 further including one or more buffering agents in combination with the statin.
11. The pharmaceutical composition as defined in Claim 3 further including an outer protective coating or finishing layer surrounding said tablet.

12. The pharmaceutical composition as defined in  
Claim 1 wherein the aspirin is in the form of enteric  
coated aspirin granules.

13. The pharmaceutical composition as defined in  
5 Claim 12 in the form of tablets or granules contained in a  
capsule.

14. The pharmaceutical composition as defined in  
Claim 12 wherein the enteric coated aspirin granules  
include a finishing overcoat, and the coated aspirin and  
10 statin are in the form of a tablet or capsule.

15. The pharmaceutical composition as defined in  
Claim 14 wherein the coated aspirin granules and statin  
granules are encapsulated separately into the same capsule  
shells.

16. The pharmaceutical composition as defined in  
Claim 2 wherein the aspirin is in the form of enteric  
coated granules of aspirin and the statin is in the form of  
enteric coated granules of statin, in the form of  
compressed tablets or capsules.

17. The pharmaceutical composition as defined in  
Claim 1 in the form of a tablet or capsule containing both  
aspirin granules and statin granules.

18. The pharmaceutical composition as defined in  
Claim 17 wherein the statin is in the form of enteric  
25 coated statin granules.

19. The pharmaceutical composition as defined in  
Claim 17 wherein the statin granules include an outer  
protective coating to minimize interaction with aspirin.

20. The pharmaceutical composition as defined in  
30 Claim 1 in the form of a bilayered tablet which comprises a  
first layer comprising aspirin granules and one or more  
excipients, and a second layer comprising a statin and one  
or more buffering compounds and one or more excipients.

21. The pharmaceutical composition as defined in  
35 Claim 20 wherein the first layer comprises aspirin  
granules, one or more bulking agents and optionally a  
lubricant, and the second layer comprises a statin.

optionally a wet granulating agent, one or more buffering compounds selected from the group consisting of calcium carbonate, magnesium oxide, magnesium carbonate and mixtures thereof, and optionally magnesium stearate.

5        22. The pharmaceutical composition as defined in Claim 20 further including an outer protective coating surrounding said bilayered tablet.

23. The pharmaceutical composition as defined in Claim 1 further including an antioxidant.

10        24. The pharmaceutical composition as defined in Claim 23 wherein the antioxidant is vitamin C and/or vitamin E.

15        25. A method for lowering serum cholesterol or preventing or inhibiting or treating atherosclerosis or reducing risk of or treating a cardiovascular event or disease, coronary artery disease or cerebrovascular disease, which comprises administering to a patient in need of treatment a therapeutically effective amount of a pharmaceutical composition comprising a combination of a 20 statin cholesterol lowering agent and aspirin in a single dosage form, which dosage form minimizes interaction between the statin and aspirin, and reduces the side effects of aspirin.

25        26. The method as defined in Claim 25 wherein the statin employed is pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/05111

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 9/16, 9/24, 9/26, 9/30, 9/48, 9/52

US CL : 424/452, 458, 459, 470, 472, 474, 489, 490

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/452, 458, 459, 470, 472, 474, 489, 490

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4,642,233 A (URQUHART et al) 10 February 1987, abstract, column 6, line 25.	1-14, 17-19

<input type="checkbox"/>	Further documents are listed in the continuation of Box C.	<input type="checkbox"/>	See patent family annex.
*A*	Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A*	document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B*	earlier documents published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L*	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	*A*	document member of the same patent family
*O*	document referring to an oral disclosure, use, exhibition or other means		
*P*	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search  07 JUNE 1999	Date of mailing of the international search report  07 JUL 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer  <i>Lawrence T</i> Edward J. Webman Telephone No. (703) 308-1234

1/9/1

15414608

Basic Patent (No,Kind,Date): WO 9947123 A1 19990923

## PATENT FAMILY:

## AUSTRALIA (AU)

Patent (No,Kind,Date): AU 9929015 A1 19991011

PHARMACEUTICAL COMPOSITION CONTAINING A STATIN AND ASPIRIN (English)

Patent Assignee: SQUIBB BRISTOL MYERS CO

Author (Inventor): ULLAH ISMAT; JAIN NEMICHAND B

Priority (No,Kind,Date): US 40794 A 19980318; WO 99US5111 W 19990309

Applic (No,Kind,Date): AU 9929015 A 19990309

IPC: \* A61K-009/16; A61K-009/24; A61K-009/26; A61K-009/30; A61K-009/48 ; A61K-009/52

CA Abstract No: \* 131(17)233578H

Derwent WPI Acc No: \* C 99-561835

Language of Document: English

## BRAZIL (BR)

Patent (No,Kind,Date): BR 9908690 A 20001205

COMPOSICAO FARMACEUTICA CONTENDO ESTATINA E ASPIRINA (Portuguese)

Patent Assignee: SQUIBB BRISTOL MYERS CO (US)

Author (Inventor): ULLAH ISMAT; JAIN NEMICHAND B

Priority (No,Kind,Date): US 40794 A 19980318; WO 99US5111 W 19990309

Applic (No,Kind,Date): BR 99U8690 A 19990309

IPC: \* A61K-009/16; A61K-009/24; A61K-009/26; A61K-009/30; A61K-009/48 ; A61K-009/52

CA Abstract No: \* 131(17)233578H

Derwent WPI Acc No: \* C 99-561835

Language of Document: Portuguese

## CANADA (CA)

Legal Status (No,Type,Date,Code,Text):

CA 2324283 P 20000915 CA REFW CORRESPONDS TO PCT APPLICATION (ENTSPRICHT PCT ANMELDUNG)  
WO 9947123 P

## EUROPEAN PATENT OFFICE (EP)

Patent (No,Kind,Date): EP 1071403 A1 20010131

PHARMACEUTICAL COMPOSITION CONTAINING A STATIN AND ASPIRIN (English; French; German)

Patent Assignee: SQUIBB BRISTOL MYERS CO (US)

Author (Inventor): ULLAH ISMAT (US); JAIN NEMICHAND B (US)

Priority (No,Kind,Date): WO 99US5111 W 19990309; US 40794 A 19980318

Applic (No,Kind,Date): EP 99909930 A 19990309

Designated States: (National) AT; BE; CH; CY; DE; DK; ES; FI; FR; GB; GR; IE; IT; LI; LU; MC; NL; PT; SE

IPC: \* A61K-009/16; A61K-009/24; A61K-009/26; A61K-009/30; A61K-009/48 ; A61K-009/52

CA Abstract No: \* 131(17)233578H

Derwent WPI Acc No: \* C 99-561835

Language of Document: English

## EUROPEAN PATENT OFFICE (EP)

Legal Status (No,Type,Date,Code,Text):

EP 1071403 P 19980318 EP AA PRIORITY (PATENT APPLICATION) (PRIORITAET (PATENTANMELDUNG))  
US 40794 A 19980318

EP 1071403 P 19990309 EP AA PCT-APPLICATION  
 (PCT-ANMELDUNG)  
 WO 99US5111 W 19990309  
 EP 1071403 P 19990309 EP AE EP-APPLICATION  
 (EUROPAEISCHE ANMELDUNG)  
 EP 99909930 A 19990309  
 EP 1071403 P 20010131 EP AK DESIGNATED CONTRACTING  
 STATES IN AN APPLICATION WITH SEARCH REPORT:  
 (IN EINER ANMELDUNG BENANNTTE VERTRAGSSTAATEN)  
 AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
 MC NL PT SE  
 EP 1071403 P 20010131 EP A1 PUBLICATION OF APPLICATION  
 WITH SEARCH REPORT (VEROEFFENTLICHUNG DER  
 ANMELDUNG MIT RECHERCHENBERICHT)  
 EP 1071403 P 20010131 EP 17P REQUEST FOR EXAMINATION  
 FILED (PRUEFUNGSANTRAG GESTELLT)  
 20001016

WORLD INTELLECTUAL PROPERTY ORGANIZATION, PCT (WO)

Patent (No,Kind,Date): WO 9947123 A1 19990923  
 PHARMACEUTICAL COMPOSITION CONTAINING A STATIN AND ASPIRIN (English)  
 Patent Assignee: SQUIBB BRISTOL MYERS CO (US)  
 Author (Inventor): ULLAH ISMAT; JAIN NEMICHAND B  
 Priority (No,Kind,Date): US 40794 A 19980318  
 Applic (No,Kind,Date): WO 99US5111 A 19990309  
 Designated States: (National) AL; AM; AT; AU; AZ; BA; BB; BG; BR; BY;  
 CA; CH; CN; CU; CZ; DE; DK; EE; ES; FI; GB; GE; GH; GM; HU; ID; IL;  
 IS; JP; KE; KG; KP; KR; KZ; LC; LK; LR; LS; LT; LU; LV; MD; MG; MK;  
 MN; MW; MX; NO; NZ; PL; PT; RO; RU; SD; SE; SG; SI; SK; SL; TJ; TM;  
 TR; TT; UA; UG; UZ; VN; YU; ZW (Regional) GH; GM; KE; LS; MW; SD;  
 SL; SZ; UG; ZW; AM; AZ; BY; KG; KZ; MD; RU; TJ; TM; AT; BE; CH; CY;  
 DE; DK; ES; FI; FR; GB; GR; IE; IT; LU; MC; NL; PT; SE; BF; BJ; CF;  
 CG; CI; CM; GA; GN; GW; ML; MR; NE; SN; TD; TG

Filing Details: WO 100000 With international search report  
 IPC: \* A61K-009/16; A61K-009/24; A61K-009/26; A61K-009/30; A61K-009/48  
 ; A61K-009/52

CA Abstract No: \* 131(17)233578H; 131(17)233578H

Derwent WPI Acc No: \* C 99-561835; C 99-561835

Language of Document: English

WORLD INTELLECTUAL PROPERTY ORGANIZATION, PCT (WO)

Legal Status (No,Type,Date,Code,Text):  
 WO 9947123 P 19980318 WO AA PRIORITY (PATENT)  
 US 40794 A 19980318  
 WO 9947123 P 19990309 WO AE APPLICATION DATA (APPL.  
 DATA)  
 WO 99US5111 A 19990309  
 WO 9947123 P 19990923 WO AK DESIGNATED STATES CITED IN A  
 PUBLISHED APPLICATION WITH SEARCH REPORT  
 (DESIGNATED STATES CITED IN A PUBLISHED APPL.  
 WITH SEARCH REPORT)  
 AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ  
 DE DK EE ES FI GB GE GH GM HU ID IL IS JP KE  
 KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN  
 MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ  
 TM TR TT UA UG UZ VN YU ZW  
 WO 9947123 P 19990923 WO AL DESIGNATED COUNTRIES FOR  
 REGIONAL PATENTS CITED IN A PUBLISHED  
 APPLICATION WITH SEARCH REPORT (DESIGNATED  
 COUNTRIES FOR REGIONAL PATENTS CITED IN A  
 PUBLISHED APPL. WITH SEARCH REPORT)  
 GH GM KE LS MW SD SL SZ UG ZW AM AZ BY KG KZ

MD RU TJ TM AT BE CH CY DE DK ES FI FR GB GR  
IE IT LU MC NL PT SE BF BJ CF CG CI CM GA CN  
GW ML MR NE SN TD TG

WO 9947123 P 19990923 WO A1 PUBLICATION OF THE  
INTERNATIONAL APPLICATION WITH THE  
INTERNATIONAL SEARCH REPORT (PUB. OF THE  
INTERNATIONAL APPL. WITH THE INTERNATIONAL  
SEARCH REPORT)  
WO 9947123 P 19991117 WO 121 EP: PCT APP. ART. 158 (1)  
(EP: PCT ANM. ART. 158 (1))  
WO 9947123 P 20000106 WO DFPE REQUEST FOR PRELIMINARY  
EXAMINATION FILED PRIOR TO EXPIRATION OF 19TH  
MONTH FROM PRIORITY DATE  
WO 9947123 P 20000915 WO ENP ENTRY INTO THE NATIONAL  
PHASE IN:  
CA 2324283 AA  
WO 9947123 P 20010118 DE 8642/REG IMPACT ABOLISHED FOR DE  
(WIRKUNG WEGGEFALLEN FUER DE)

Inpadoc Fam. & Legal Stat (Dialog & File 345); (c) 2001 EPO. All rights reserved.

© 2001 The Dialog Corporation